

Una técnica alternativa para el cariotipado mitótico en saltamontes: bandeo C y Fluorescente en *Adimantus ornatissimus* (Orthoptera: Acrididae)

An alternative technique for mitotic grasshoppers karyotyping: Fluorescent and C-banding in *Adimantus ornatissimus* (Orthoptera: Acrididae)

Elio R. Castillo, Alberto Taffarel, Dardo A. Martí

Resumen

Los ortópteros son el material de preferencia para estudios citogenéticos debido al tamaño de sus cromosomas y a su bajo número diploide, sin embargo existen dificultades para obtener cromosomas mitóticos de alta calidad para tinciones diferenciales, siguiendo un protocolo fácil y reproducible. Con la intención de optimizar la calidad de los cromosomas mitóticos y minimizar el material no cromosómico en las preparaciones, se han utilizado diferentes procedimientos y tejidos. Los preparados mitóticos se realizan macerando ciegos gástricos y posteriormente se utiliza hielo seco, CO₂ a alta presión o nitrógeno líquido para remover el cubreobjetos. En el presente trabajo describimos una técnica que optimiza la visualización y la caracterización morfológica de los cromosomas mitóticos y facilita la aplicación de los procedimientos de tinción diferencial. Como material citológico de referencia analizamos a *Adimantus ornatissimus* (Burmeister 1838), la especie posee un $2n=23/24$ ♂/♀, y un sistema de determinación sexual XO/XX.

Palabras clave: ciegos gástricos; mitosis; cromosomas; Orthoptera; técnica de frotis.

Abstract

Orthopterans are a preferred material for cytogenetic studies because of their large chromosome size and low diploid number; however a difficulty is to maximize a high quality mitotic chromosomes obtention and the choice of an easy and reproducible protocol. Different procedures and tissues were used to optimize mitotic chromosome observation with conventional staining, preserving, as much as possible chromosome morphology, and minimizing the non-chromosomal background material. Grasshopper mitotic metaphases preparations are usually obtained from squashed female gastric caeca and posterior exposure of slides to dry ice, CO₂ flow, or liquid nitrogen to remove coverslips. Grasshopper embryos are also used for cytological analyses. As a reference cytological material, we analyzed *Adimantus ornatissimus* (Burmeister 1838) chromosomes. This specie showed $2n=23/24$ ♂/♀ and a XO/XX sex chromosome system. We describe the frotis technique which optimizes the visualization and characterization of mitotic chromosome morphology and facilitates differential staining procedures.

Key words: gastric caecum, mitosis, chromosomes, Orthoptera, frotis technique.

Introduction

Species of Orthoptera are distributed over a wide range of habitats. However, as it happens in other groups of animals, a significant amount of species included in the order are confined to the Neotropical region, and only a few to cold latitudes [10]. Most of Neotropical grasshopper diversity is represented by Acridoidea [29], which include the family Acrididae, the most widely studied at the cytogenetic level [10, 29].

Grasshoppers are an ideal material for basic and population cytogenetic studies because of large chromosome size, low chromosome number, and exceptionally clear

meiosis [10]. Used as experimental material, they contributed significantly to the knowledge of chromosome behavior in mitosis and meiosis including pairing, synapsis, chiasma formation and recombination, and to the study of numerical and structural mutations and chromosomal polymorphisms [3, 5, 10, 13, 14, 19, 22, 26].

Since many decades Acridoids cytogenetic analyses have been performed and based on male individuals due the easiness of meiotic chromosome preparations. On the other hand, female meiosis presents many difficulties for chromosome study [8, 15].

An additional difficulty is to choose an adequate protocol to maximize the obtention of high quality mi-

totic chromosomes. Different procedures and tissues are used to optimize mitotic chromosome observation with conventional staining preserving, as much as possible, chromosome morphology, and minimizing the background material, producing slides with well-spread and complete metaphases. Moreover, mitotic preparations have to be usually manipulated for differential staining, including fluorochromes [25]. Generally, mitotic chromosome preparations of grasshoppers are obtained by squashing female gastric caeca after incubation with a mitotic poison [12, 17, 18, 24]. Slides are then exposed to dry ice, CO₂ flow or liquid nitrogen immersion to remove the coverslip [7, 21]. Mitotic cytological preparations are also obtained through squashing of females ovarioles [7, 17, 20].

Grasshopper embryos are also cultured for mitotic chromosome obtention [1, 11, 25]. However, additional care must be taken when the protocol is developed, like extractions of embryos from the pods and in the first steps of breeding [1, 11].

We present an improved, alternative and easy technique that facilitates and optimizes the visualization and characterization of mitotic chromosome morphology in grasshoppers. The protocol was adapted from a technique used in vertebrates from intestinal tissue [23], and modified for orthopteran species in our laboratory (see below) mainly using less works solutions and replacing the intestinal suspension dropping by the frotis procedure.

Materials and methods

Test species. Eight adult females of *Adimantus ornatissimus* (Burmeister 1838) (Orthoptera, Acrididae) were studied at eight localities of Misiones province, Argentina (Fig. 1).

Mitotic preparations

Mitotic chromosomes were obtained from intestinal cells of gastric caecum through a new air-drying technique adapted by us to orthopteran, from a protocol used in vertebrates. The same is as follow:

1. Females were injected at the level of the fifth abdominal segment with 0.05% colchicine in insect saline (CINa 0.9%) for 4-5 hours.
2. After colchicine exposure, gastric caeca were dissected and immersed in a KCl 0.75% hypotonic solution for 45-55 minutes.
3. Before definitive fixation in 3:1 methanol: glacial acetic acid, tissues were passed through a battery of fixative solution to remove hypotonic solution's remains, and then stored at 4°C.
4. A small portion of gastric caecum is ruptured in 45% acetic acid. The final solution is resuspended twice using a Pasteur pipette during a period of 12 to 15 minutes.
5. Chromosomal preparations have to be performed under an airfoil chemical fume hood, equipped with an air extractor, to prevent vapor inhalation.



Figure 1. Female habitus of *Adimantus ornatissimus* from Moconá, San Pedro, Misiones, Argentina.

6. A drop of the suspension is then placed on one end of a warm slide, from which the material is extended towards the opposite side, with the aid of another glass slide (frotis). The last step is repeated from one side of the slide to the other until no liquid is left.
7. Finally, slides are air-dried at room temperature.

Staining. For solid staining phosphate buffered 5% Giemsa is used. For sequential staining, C banding [27], and C-DAPI banding [2] destaining must be performed. After observation and photomicrography of mitotic metaphases, immersion oil is removed from slides by three successive immersions in pure Xylene, washed in distilled water, and air dried at room temperature. Then, for Giemsa removal, slides are immersed in 3:1 methanol:glacial acetic acid. Double staining with CMA3/DAPI was done as described by Schweizer *et al* (1983) [28].

Results and discussion

The first record of *A. ornatissimus* males and females chromosome number was proposed by Mesa *et al* (1982) [16] but cell draw or photomicrography has never been showed or described. We observed a $2n=23$ diploid number in males [16], and our results agree with the female chromosome set which shows $2n=24$ ($FN=24$) telocentric chromosomes (Fig. 2 a). This karyotypic constitution is common in females of Acrididae [5, 16]. Based on chromosome size measurements three distinct groups were distinguished: three pairs of large chromosomes (L1-L3), five medium sized (M4-M8), three small (S9-S11), and the sex chromosomes which is similar in size to M4.

In most cytogenetic studies, using grasshoppers as experimental organisms, males are more frequently studied, because testes dissection is easy (with basic anatomical knowledge), and a lot of meiotic chromosome preparations per male individual can be obtained, with an elevated percentage of cells in different meiotic phases [9]. Male meiotic preparations don't present major difficulties and reveal relevant details of chromosome morphology, bivalents configurations, different condensation patterns (as happens in sex chromosomes at post-pachytene phases), chromosome segregation, nucleolar cycle, and also recombination sites, represented by chiasmata.

A radically different situation occurs with female meiosis, where a systematic and laborious protocol needs to be followed, from the initial capture of females through the correct dissection of the egg, to the obtention of a single first metaphase per egg. Gravid females have to be fed and oviposition has to be stimulated. When oviposition is going to take place, eggs are extracted and the micropilar region of the primary oocyte is dissected out and gently squashed [15]. The yield of the technique is extremely low.

On the other hand, mitotic chromosome information is relevant to determine the chromosome number and cytolo-

gical characteristics like constitutive heterochromatin and NORs [4, 6, 25]. Thus, in females, mitosis is the main source of chromosomal information and to that end, a number of different tissues (e.g. caeca, ovaries and embryos) and techniques are usually employed [4, 6, 25]. However, some procedures like embryo culture are laborious, requiring a lot of time and cares [4]; others do not yield good quality chromosomes.

The technique described in this paper maximizes the percentage (ca. 90%) of well-spread, high quality and complete metaphases in each preparation. In the same way, slide preparation involves less time, equipment, cost and space in comparison with other protocols, for example, when the coverslips have to be removed with ice dry, liquid nitrogen or CO_2 [7, 21]. One disadvantage of liquid nitrogen, dry ice or CO_2 procedures is the lost of material when the coverslips is removed (personal observation). In most of the cases, part of the preparation remains in the slide and part in the coverslips. Although use of siliconized coverslips prevents the loss of material in simple chromosome squashes, the protocol requires more time compared with our technique. Another difficulty of this protocol is the distribution of chromosomes in different focal planes (personal observation).

With the alternative frotis procedure the totality of intestinal material and also the complete slide surface is used. In standard squash caeca preparations, the material is confined to a limited area. Our procedure allows complete spread cells with chromosomes in one focal plane (Fig. 2 a, b, c, d, e). As happens with other procedures, mitotic metaphases treated with Giemsa, can be used in differential staining techniques after slides decoloration took place (Fig. 2 c, d, e).

The technique proposed in this paper for orthopteran mitotic chromosome preparations combines high quality, perfect reproducibility of the results and processing speed, with minimum laboratory conditions dispensing the use of dry ice or mechanical procedures, normally used in coverslips removal.

Acknowledgements

ERC is PhD candidate at Universidad Nacional de Córdoba (Doctorado en Ciencias Biológicas) and also acknowledges Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Comité Ejecutivo de Desarrollo e Innovación Tecnológica (CEDIT). DAM acknowledges the continuous support of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). This work was partially financed through grant PICTO 37035 FONCyT to DAM.




Figure 2. Female mitotic chromosomes of *Adimantus ornatissimus*. (a) Karyotype with Giemsa staining, (b) C-banding; constitutive heterochromatin patterns are dark stained and (c) sequential C-DAPI staining. DAPI/CMA3 staining, arrowheads signaling negative DAPI heterochromatic blocks in (d) and chromomycin-positive sites in (e). Bar = 10 µm.

Bibliography

1. Araujo, D. Rheims, C.A. Brescovit, A.D., Cella, D.M. *Extreme degree of chromosome number variability in species of the spider genus Scytodes (Araneae, Haplogynae, Scytodidae)*. J. Zool Syst Evol Res 46(2): 89-95, 2008.
2. Barros e Silva, A.E., Guerra, M. *The meaning of DAPI bands observed after C-banding and FISH procedures*. Biotech Histochem 85:115-125, 2010.
3. Bidau, C.J. *Causes of chiasma repatterning due to centric fusions*. Brazil J Genet 16 (2): 283-296, 1993.
4. Camacho, J.P.M. Viseras, E. Navas, J., Cabrero, J. *C-heterochromatin content of supernumerary chromosome segments of grasshoppers: detection of an euchromatic extra segment*. Heredity 53(1): 167-175, 1984.
5. Castillo, E.R.D. Bidau, C.J., Martí, D.A. *Neo-sex chromosome diversity in Neotropical melanopline grasshoppers (Melanoplinae, Acrididae)*. Genetica 138: 775-786, 2010.
6. Cella, D.M. Ferreira, A. *The cytogenetics of Abracris flavolineata (Orthoptera, Caelifera, Ommatolampinae, Abracrini)*. Brazil J Genet 14(2) 315-329, 1991.
7. Czaker, R. *Silver staining in transcriptionally active NOR of meiotic and mitotic cells in Acheta domesticus (L.) (Orthoptera)*. Chromosoma 68:187-193, 1978.
8. Grieco, M.L., Bidau, C.J. *Chiasma frequency and distribution in males and females of Metaleptea brevicornis adspersa (Acridinae, Acrididae) with and without B chromosomes*. Hereditas 131: 101-107, 1999.
9. Guerra, M. *Hematoxylin: a simple, multiple-use dye for chromosome analysis*. Genet Mol Biol 22(1): 77-80, 1999.

10. Hewitt, G.M. *Orthoptera. Grasshoppers and crickets*. In *Animal Cytogenetics 3. Insecta 1*. (B. John, ed.) Gebrüder Borntraeger, Berlin-Stuttgart. 170p, 1979.
11. John, B., King, M. *Pseudoterminalisation, terminalisation and non-chiasmate modes of terminal association*. *Chromosoma* 92: 89-99, 1985.
12. Kayano, H. *Accumulation of B chromosomes in the germ line of Locusta migratoria*. *Heredity* 27: 119-123, 1971.
13. King, M. *Chromosomal rearrangements, speciation and the theoretical approach*. *Heredity* 59: 1-6, 1987.
14. King, M. *Species Evolution. The Role of chromosome change*. Cambridge University Press. Cambridge 364p. 1993.
15. Martí, D.A., Bidau, C.J. *Male and female meiosis in a natural population of Dichroplus pratensis (Acrididae) polymorphic for Robertsonian translocation: A study of chiasma frequency and distribution*. *Hereditas* 123: 227-235, 1995.
16. Mesa, A., Ferreira, A., Carbonell, C.S. *Cariología de los acridoideos neotropicales: estado actual de su conocimiento y nuevas contribuciones*. *Anns. Soc. ent. Fr.*, 18, 4: 507-526, 1982.
17. Nankivell, R.N. *A terminal association of two pericentric inversions in first metaphase cells of the Australian grasshopper Austroicetes interioris (Acrididae)*. *Chromosoma* 22, 42-68. 1967.
18. Nur, U. *Mitotic instability leading to an accumulation of B-chromosomes in grasshoppers*. *Chromosoma* 27: 1-19, 1969.
19. Perry, P.E., Jones, G.H. *Male and female meiosis in Grasshoppers. I. Stethophyma grossum*. *Chromosoma* 47: 227-236, 1974.
20. Rocha, M.F., Souza, M.J., Moura, R.C. *Karyotype analysis, constitutive heterochromatin and NOR distribution in five grasshopper species of the subfamily Leptysminae (Acrididae)*. *Caryologia* 57:107-116, 2004.
21. Rodríguez Iñigo, E., Mason, P.L., Rufas, J.S., García de la Vega, C. *Effects of supernumerary heterochromatin on chiasma formation and chromosome segregation in Dociostaurus genei (Orthoptera)*, *Heredity*, 80: 353-360, 1998.
22. Sanchez Rufas, J. *Contribución de las técnicas de impregnación argéntica al estudio de los cromosomas de Ortópteros*. In: Gosálvez, J., López Fernández, C. and García de la Vega, C. (eds) *Orthoptera*, 1: 227-255, 1985. Fundación Ramón Areces, Madrid.
23. Schmid, M., Olert, J., Klett, C. *Chromosome banding in Amphibia III. Sex chromosomes in Triturus*. *Chromosoma* 71:29-55, 1979.
24. Souza, M.J., Haver, P.R.O., Melo, N.F. *Karyotype, C- and fluorescence banding patterns, NORs location and FISH in the grasshopper Xestotrachelus robustus (Romaleidae)*. *Caryologia* 56: 261-267, 2003.
25. Souza, M.J., Melo, N.F. *Chromosome study in Schistocerca (Orthoptera, Acrididae, Cyrtacanthacridinae): Karyotypes and distribution patterns of constitutive heterochromatin and nucleolus organizer regions (NORs)*. *Genet Mol Biol* 30(1): 54-59, 2007.
26. Suja, J.A. Antonio, C. González-García, J.M., Rufas, J.S. *Supernumerary heterochromatic segments associated with the nucleolar chromosomes of Pyrgomorpha conica (Orthoptera) contain methylated rDNA sequences*. *Chromosoma* 102: 491-499, 1993.
27. Sumner, A.T. *A simple technic for demonstrating centromeric heterochromatin*. *Exp Cell Res* 75: 304- 306, 1972.
28. Schweizer, D. Mendelak, M. White, M.J.D., Contreras, N. *Cytogenetics of the parthenogenetic grasshopper Warramaba virgo and its bisexual relatives. X. Patterns of fluorescent banding*. *Chromosoma* 88: 227-236, 1983.
29. White, M.J.D. *Animal cytology and evolution*. Cambridge University Press. Cambridge, 3rd ed, 1973.

Recibido: 10/03/11

Aprobado: 04/03/12

• Elio Rodrigo Castillo^{1,2,3}

Licenciado en Genética de la UNaM. Becario de CONICET-CEDIT y realiza el Doctorado Ciencias Biológicas en la Universidad Nacional de Córdoba. Desempeña actividades de docencia en la Cátedra de Genética Evolutiva de la F.C.E.Q.yN. (UNaM) con el cargo de Ayudante de Primera. Correo electrónico: castillo.eliorodrigo@gmail.com

• Alberto Taffarel¹

Se encuentra desarrollando su tesis de grado de la carrera Licenciatura en Genética, bajo la dirección del Dr. Dardo Martí, en el Laboratorio de Genética Evolutiva de la UNaM. Correo electrónico: radova@gmail.com

• Dardo Andrea Martí^{1,2}

Dr. en Ciencias Biológicas de la Universidad Nacional de Córdoba. El Dr. Martí es Investigador Adjunto de CONICET. Asimismo, se encuentra bajo su dirección el Laboratorio de Genética Evolutiva de la F.C.E.Q.yN. (UNaM) y es docente de la cátedra de Genética Evolutiva desempeñando el cargo de Profesor Adjunto Regular. Correo electrónico: darmarti@fceqyn.unam.edu.ar

1. Universidad Nacional de Misiones, Facultad de Ciencias Exactas, Químicas y Naturales, Laboratorio de Genética Evolutiva. Félix de Azara 1552. CP. 3300 Posadas, Misiones, Argentina.

2. Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). Rivadavia 1917, 1033. Ciudad Autónoma de Buenos Aires. Argentina.

3. Comité Ejecutivo de Desarrollo e Innovación Tecnológica (CEDIT).